

# Quadrilaterals

## Multiple Choice Questions (MCQs)

1. How many angles are there in a quadrilateral?

- (a) 4 (b) 2 (c) 1 (d) 3

2. The three consecutive angles of a quadrilateral are  $70^\circ$ ,  $120^\circ$  and  $50^\circ$ . The fourth angle of the quadrilateral is

- (a)  $45^\circ$  (b)  $60^\circ$  (c)  $120^\circ$  (d)  $30^\circ$

3. If the sum of angles of a triangle is  $X$  and the sum of the angles of a quadrilateral is  $Y$ , then

- (a)  $X = 2Y$  (b)  $2X = Y$   
 (c)  $X = Y$  (d)  $X + Y = 360^\circ$

4. One of the angles of a quadrilateral is  $90^\circ$  and the remaining three angles are in the ratio  $2:3:4$ . Find the largest angle of the quadrilateral.

- (a)  $120^\circ$  (b)  $90^\circ$  (c)  $140^\circ$  (d)  $100^\circ$

5. In the figure,  $ABCD$  is a quadrilateral whose sides  $AB$ ,  $BC$ ,  $CD$  and  $DA$  are produced in order to  $P$ ,  $Q$ ,  $R$  and  $S$ . Then  $x + y + z + t$  is equal to

- (a)  $180^\circ$  (b)  $360^\circ$  (c)  $380^\circ$  (d)  $270^\circ$

6. If only one pair of opposite sides of a quadrilateral are parallel, then the quadrilateral is a

- (a) Parallelogram (b) Trapezium  
 (c) Rhombus (d) Rectangle

7. A blackboard is in the shape of a

- (a) Parallelogram (b) Rhombus  
 (c) Rectangle (d) Kite

8. The angle between the diagonals of a rhombus is

- (a)  $45^\circ$  (b)  $90^\circ$  (c)  $30^\circ$  (d)  $60^\circ$

9. A quadrilateral whose all the four sides and all the four angles are equal is called a

- (a) Rectangle (b) Rhombus  
 (c) Square (d) Parallelogram

10. Which of the following is not true?


- (a) The diagonals of a rectangle are equal.  
 (b) Diagonals of a square are equal.

(c) Diagonals of a parallelogram are not always equal.

(d) Diagonals of a kite are equal.

11. In the adjoining figure,  $ABCD$  is a square. A line segment  $DX$  cuts the side  $BC$  at  $X$  and the diagonal  $AC$  at  $O$  such that  $\angle COD = 105^\circ$  and  $\angle OXC = x^\circ$ . Find the value of  $x$ .

- (a)  $75^\circ$  (b)  $80^\circ$  (c)  $60^\circ$  (d)  $45^\circ$



12. If angles  $A$ ,  $B$ ,  $C$  and  $D$  of the quadrilateral  $ABCD$ , taken in order, are in the ratio  $3:7:6:4$ , then  $ABCD$  is a


- (a) rhombus (b) parallelogram  
 (c) trapezium (d) kite

13. In a parallelogram  $ABCD$ , if  $\angle A = 75^\circ$ , then the measure of  $\angle B$  is

- (a)  $10^\circ$  (b)  $20^\circ$  (c)  $105^\circ$  (d)  $90^\circ$

14. In parallelogram  $ABCD$ ,  $\angle DAB = 70^\circ$ ,  $\angle DBC = 70^\circ$ , then  $\angle CDB$  is equal to


- (a)  $40^\circ$   
 (b)  $60^\circ$   
 (c)  $70^\circ$   
 (d)  $30^\circ$



15. In the given figure,  $ABCD$  is a parallelogram.  $E$  and  $F$  are points on opposite sides  $AD$  and  $BC$  respectively, such

that  $ED = \frac{1}{2}AD$  and  $BF = \frac{1}{3}BC$ . If  $\angle ADF = 60^\circ$ , then find  $\angle BFD$ .

- (a)  $120^\circ$  (b)  $130^\circ$  (c)  $125^\circ$  (d)  $115^\circ$




16. Two angles of a quadrilateral are  $55^\circ$  and  $65^\circ$ . The other two angles are in the ratio  $3:5$ . The two angles are

- (a)  $100^\circ$ ,  $110^\circ$  (b)  $85^\circ$ ,  $125^\circ$   
 (c)  $100^\circ$ ,  $120^\circ$  (d)  $90^\circ$ ,  $150^\circ$

17. In a quadrilateral  $ABCD$ , diagonals bisect each other at right angle. Also,  $AB = BC = AD = 5$  cm, then find the length of  $CD$ .

- (a) 5 cm (b) 4 cm (c) 2 cm (d) 6 cm

18. In the given figure,  $ABCD$  is a parallelogram, what is the sum of the angles  $x$ ,  $y$  and  $z$ ?




- (a)  $180^\circ$  (b)  $45^\circ$  (c)  $60^\circ$  (d)  $90^\circ$

19. If a pair of opposite sides of a quadrilateral is equal and parallel, then the quadrilateral is a

- (a) parallelogram (b) rectangle  
(c) rhombus (d) square

20. In  $\triangle ABC$ ,  $EF \parallel BC$ ,  $F$  is the mid-point of  $AC$  and  $AE = 3.5$  cm. Then  $AB$  is equal to



- (a) 7 cm  
(b) 5 cm  
(c) 5.5 cm  
(d) 4.5 cm

21. The triangle formed by joining the mid-points of the sides of an equilateral triangle is

- (a) scalene (b) right angled  
(c) equilateral (d) isosceles


22. The four triangles formed by joining the mid-points of the sides of a triangle are

- (a) congruent to each other  
(b) non-congruent to each other  
(c) always right angled triangle  
(d) can't be determined

23. If  $M$  and  $N$  are the mid-points of non parallel sides of a trapezium  $PQRS$ , then which of the following conditions is/are true?

- (a)  $MN \parallel PQ$   
(b)  $MN = \frac{1}{2} (PQ + RS)$   
(c)  $MN = \frac{1}{2} (PQ - RS)$   
(d) Both (a) and (b)

24. In the given figure,  $ABCD$  is a rhombus. If  $\angle A = 70^\circ$ , then  $\angle CDB$  is equal to



- (a)  $65^\circ$   
(b)  $55^\circ$   
(c)  $75^\circ$   
(d)  $80^\circ$


25. Two adjacent angles of a parallelogram are  $(2x + 25)^\circ$  and  $(3x - 5)^\circ$ . The value of  $x$  is

- (a) 28 (b) 32 (c) 36 (d) 42

26. In a quadrilateral  $STAR$ , if  $\angle S = 120^\circ$ , and  $\angle T : \angle A : \angle R = 5 : 3 : 7$ , then measure of  $\angle R =$

- (a)  $112^\circ$  (b)  $120^\circ$   
(c)  $110^\circ$  (d) None of these

27. In figure,  $ABCD$  is a trapezium. Find the values of  $x$  and  $y$ .



- (a)  $x = 50^\circ, y = 80^\circ$   
(b)  $x = 50^\circ, y = 88^\circ$   
(c)  $x = 80^\circ, y = 50^\circ$   
(d) None of these

28. In a quadrilateral  $ABCD$ ,  $\angle A + \angle C$  is 2 times  $\angle B + \angle D$ . If  $\angle A = 140^\circ$  and  $\angle D = 60^\circ$ , then  $\angle B =$

- (a)  $60^\circ$  (b)  $80^\circ$   
(c)  $120^\circ$  (d) None of these

29. The measure of all the angles of a parallelogram, if an angle is 24 less than twice the smallest angle, is

- (a)  $37^\circ, 143^\circ, 37^\circ, 143^\circ$   
(b)  $108^\circ, 72^\circ, 108^\circ, 72^\circ$   
(c)  $68^\circ, 112^\circ, 68^\circ, 112^\circ$   
(d) None of these


30. Which type of quadrilateral is formed when the angles  $A, B, C$  and  $D$  are in the ratio  $2 : 4 : 5 : 7$ ?

- (a) Rhombus (b) Square  
(c) Trapezium (d) Rectangle

31. In  $\triangle PQR$ ,  $A$  and  $B$  are respectively the mid-points of sides  $PQ$  and  $PR$ . If  $\angle PAB = 60^\circ$ , then  $\angle PQR =$


- (a)  $40^\circ$  (b)  $80^\circ$  (c)  $60^\circ$  (d)  $70^\circ$

32. Sides  $AB$  and  $CD$  of a quadrilateral  $ABCD$  are extended as in figure. Then  $a + b$  is equal to



- (a)  $x + 2y$   
(b)  $x - y$   
(c)  $x + y$   
(d)  $2x + y$

33. In the adjoining figure,  $PQRS$  is a parallelogram in which  $PQ$  is produced to  $T$  such that  $QT = PQ$ . Then,  $OQ$  is equal to



- (a)  $OS$  (b)  $OR$   
(c)  $OT$  (d) None of these

34. If consecutive sides of a parallelogram are equal, then it is necessarily a

- (a) Rectangle (b) Rhombus  
(c) Trapezium (d) None of these


35. The triangle formed by joining the mid-points of the sides of a right angled triangle is

- (a) scalene (b) isosceles  
(c) equilateral (d) right angled

## Case Based MCQs

**Case I.** Read the following passage and answer the questions from 36 to 40.

Laveena's class teacher gave students some colourful papers in the shape of quadrilaterals. She asked students to make a parallelogram from it using paper folding. Laveena made the following parallelogram.



**36.** How can a parallelogram be formed by using paper folding?

- Joining the sides of quadrilateral
- Joining the mid-points of sides of quadrilateral
- Joining the various quadrilaterals
- None of these

**37.** Which of the following is true?

- $PQ = BD$
- $PQ = \frac{1}{2}BD$
- $3PQ = BD$
- $PQ = 2BD$

**38.** Which of the following is correct combination?

- $2RS = BD$
- $RS = \frac{1}{3}BD$
- $RS = BD$
- $RS = 2BD$

**39.** Which of the following is correct?


- $SR = 2PQ$
- $PQ = SR$
- $SR = 3PQ$
- $SR = 4PQ$

**40.** Write the formula used to find the perimeter of quadrilateral PQRS.

- $PQ + QR + RS + SP$
- $PQ - QR + RS - SP$
- $\frac{PQ + QR + RS + SP}{2}$
- $\frac{PQ + QR + RS + SP}{3}$

**Case II.** Read the following passage and answer the questions from 41 to 45.

After summervacation, Manit's class teacher organised a small MCQ quiz, based on the properties of quadrilaterals.



During quiz, she asks different questions to students.

Some of the questions are listed below.

**41.** Which of the following is/are the condition(s) for ABCD to be a quadrilateral?


- The four points A, B, C and D must be distinct and co-planar.
- No three of points A, B, C and D are collinear.
- Line segments i.e., AB, BC, CD, DA intersect at their end points only.
- All of these

**42.** Which of the following is wrong condition for a quadrilateral said to be a parallelogram?

- Opposite sides are equal
- Opposite angles are equal
- Diagonal can't bisect each other
- None of these


**43.** If AX and CY are the bisectors of the angles A and C of a parallelogram ABCD, then

- $AX \parallel CY$
- $AX \parallel CD$
- $AX \parallel AB$
- None of these



**44.** ABCD and AEF G are two parallelograms. If  $\angle C = 63^\circ$ , then determine  $\angle G$ .

- $63^\circ$
- $117^\circ$
- $90^\circ$
- $120^\circ$




**45.** If angles of a quadrilateral are in ratio 3 : 5 : 5 : 7, then find all the angles.

- $54^\circ, 80^\circ, 80^\circ, 146^\circ$
- $34^\circ, 100^\circ, 100^\circ, 126^\circ$
- $54^\circ, 90^\circ, 90^\circ, 126^\circ$
- None of these

**Case III.** Read the following passage and answer the questions from 46 to 50.

Anjali and Meena were trying to prove mid point theorem.

They draw a triangle ABC, where D and E are found to be the midpoints of AB and AC respectively. DE was joined and extended to F such that  $DE = EF$  and FC is also joined.



**46.**  $\triangle ADE$  and  $\triangle CFE$  are congruent by which criterion?

- SSS
- SAS
- RHS
- ASA

47.  $\angle EFC$  is equal to which angle?  
 (a)  $\angle DAE$  (b)  $\angle EDA$  (c)  $\angle AED$  (d)  $\angle DBC$
48.  $\angle ECF$  is equal to which angle?  
 (a)  $\angle EAD$  (b)  $\angle ADE$  (c)  $\angle AED$  (d)  $\angle B$

49.  $CF$  is equal to  
 (a)  $EC$  (b)  $BE$  (c)  $BC$  (d)  $AD$
50.  $CF$  is parallel to  
 (a)  $AE$  (b)  $CE$  (c)  $BD$  (d)  $AC$

## ➡ Assertion & Reasoning Based MCQs


**Directions (Q.51 to 55) :** In these questions, a statement of Assertion is followed by a statement of Reason is given. Choose the correct answer out of the following choices :

- (a) Assertion and Reason both are correct statements and Reason is the correct explanation of Assertion.  
 (b) Assertion and Reason both are correct statements but Reason is not the correct explanation of Assertion.  
 (c) Assertion is correct statement but Reason is wrong statement.  
 (d) Assertion is wrong statement but Reason is correct statement.

**51. Assertion :** In  $\triangle ABC$ , median  $AD$  is produced to  $X$  such that  $AD = DX$ . Then  $ABXC$  is a parallelogram.

**Reason :** Diagonals of a parallelogram are perpendicular to each other.

**52. Assertion :**  $ABCD$  and  $PQRC$  are rectangles and  $Q$  is the mid-point of  $AC$ . Then  $DP = PC$ .



**Reason :** The line segment joining the mid-points of any two sides of a triangle is parallel to the third side and equal to half of it.

**53. Assertion :** Two opposite angles of a parallelogram are  $(3x - 2)^\circ$  and  $(50 - x)^\circ$ . The measure of one of the angle is  $37^\circ$ .

**Reason :** Opposite angles of a parallelogram are equal.

**54. Assertion :**  $ABCD$  is a square.  $AC$  and  $BD$  intersect at  $O$ . The measure of  $\angle AOB = 90^\circ$ .

**Reason :** Diagonals of a square bisect each other at right angles.

**55. Assertion :** In  $\triangle ABC$ ,  $E$  and  $F$  are the midpoints of  $AC$  and  $AB$  respectively. The altitude  $AP$  at  $BC$  intersects  $FE$  at  $Q$ . Then,  $AQ = QP$ .


**Reason :** If  $Q$  is the midpoint of  $AP$ , then  $AQ = QP$ .

## SUBJECTIVE TYPE QUESTIONS

### ➡ Very Short Answer Type Questions (VSA)


1. Two consecutive angles of a parallelogram are  $(x + 60^\circ)$  and  $(2x + 30^\circ)$ . What special name can you give to this parallelogram?

2. In the given figure,  $PQRS$  is a parallelogram in which  $\angle PSR = 125^\circ$ . Find the measure of  $\angle RQT$ .




3. Can the angles  $110^\circ$ ,  $80^\circ$ ,  $70^\circ$  and  $95^\circ$  be the angles of a quadrilateral? Why or why not?

4. In the figure, it is given that  $QLMN$  and  $NLRM$  are parallelograms. Can you say that  $QL = LR$ ? Why or why not?




5.  $ABCD$  is a parallelogram in which  $\angle A = 78^\circ$ . Compute  $\angle B$ ,  $\angle C$  and  $\angle D$ .

6. In the given figure,  $ABCD$  is a parallelogram in which  $\angle DAB = 60^\circ$  and  $\angle DBC = 55^\circ$ . Compute  $\angle CDB$  and  $\angle ADB$ .



7. In the given figure,  $AB = AC$  and  $CP \parallel BA$  and  $AP$  is the bisector of exterior  $\angle CAD$  of  $\triangle ABC$ . Prove that  $\angle PAC = \angle BCA$  and  $ABCP$  is a parallelogram.



8. If one angle of a rhombus is a right angle, then it is necessarily a \_\_\_\_\_ .
9. In a rhombus  $ABCD$ , if  $\angle A = 60^\circ$ , then find the sum of  $\angle A$  and  $\angle C$ .


10.  $ABCD$  is a trapezium in which  $AB \parallel DC$  and  $\angle A = \angle B = 45^\circ$ . Find angles  $C$  and  $D$  of the trapezium.

## ➡ Short Answer Type Questions (SA-I)

11. In a quadrilateral  $ABCD$ ,  $CO$  and  $DO$  are the bisectors of  $\angle C$  and  $\angle D$  respectively.

Prove that  $\angle COD = \frac{1}{2}(\angle A + \angle B)$ .

12. In the given parallelogram  $ABCD$ , the sum of any two consecutive angles is  $180^\circ$  and opposite angles are equal. Find the value of  $\angle A$ .




13. Diagonals of a quadrilateral  $ABCD$  bisect each other.  $\angle A = 45^\circ$  and  $\angle B = 135^\circ$ . Is it true? Justify your answer.

14.  $D$  and  $E$  are the mid-points of sides  $AB$  and  $AC$  respectively of triangle  $ABC$ . If the perimeter of  $\triangle ABC = 35$  cm, then find the perimeter of  $\triangle ADE$ .


15. In  $\triangle ABC$ ,  $AD$  is the median and  $DE \parallel AB$ , such that  $E$  is a point on  $AC$ . Prove that  $BE$  is another median.

16. In the given figure,  $M$ ,  $N$  and  $P$  are the midpoints of  $AB$ ,  $AC$  and  $BC$  respectively. If  $MN = 3$  cm,  $NP = 3.5$  cm and  $MP = 2.5$  cm, then find  $(BC + AC) - AB$ .



17. Let  $\triangle ABC$  be an isosceles triangle with  $AB = AC$  and let  $D$ ,  $E$  and  $F$  be the mid-points of  $BC$ ,  $CA$  and  $AB$  respectively. Show that  $AD \perp FE$  and  $AD$  is bisected by  $FE$ .

18. In the given rectangle  $ABCD$ ,  $\angle ABE = 30^\circ$  and  $\angle CFE = 144^\circ$ . Find the measure of  $\angle BEF$ .




19. The perimeter of a parallelogram is 30 cm. If longer side is 9.5 cm, then find the length of shorter side.

20. In a parallelogram  $ABCD$ , if  $\angle A = (3x - 20)^\circ$ ,  $\angle B = (y + 15)^\circ$  and  $\angle C = (x + 40)^\circ$ , then find  $x + y$  (in degrees).

## ➡ Short Answer Type Questions (SA-II)

21. In a parallelogram  $PQRS$ , if  $\angle QRS = 2x$ ,  $\angle PQS = 4x$  and  $\angle PSQ = 4x$ , then find the angles of the parallelogram.

22.  $l$ ,  $m$  and  $n$  are three parallel lines intersected by transversals  $p$  and  $q$  such that  $l$ ,  $m$  and  $n$  cut off equal intercepts  $AB$  and  $BC$  on  $p$  (see figure).




Show that  $l$ ,  $m$  and  $n$  cut off equal intercepts  $DE$  and  $EF$  on  $q$  also.

23. The side of a rhombus is 10 cm. The smaller diagonal is  $\frac{1}{3}$  of the greater diagonal. Find the length of the greater diagonal.

24. In  $\triangle ABC$ ,  $\angle A = 50^\circ$ ,  $\angle B = 60^\circ$  and  $\angle C = 70^\circ$ . Find the measures of the angles of the triangle formed by joining the mid-points of the sides of this triangle.

25. In given figure,  $ABCD$  is a parallelogram in which  $P$  is the midpoint of  $DC$  and  $Q$  is a point on  $AC$  such that  $CQ = \frac{1}{4}AC$ . If  $PQ$  produced




meet  $BC$  at  $R$ , then prove that  $R$  is a midpoint of  $BC$ .


26.  $ABCD$  is parallelogram.  $P$  is a point on  $AD$  such that  $AP = \frac{1}{3}AD$  and  $Q$  is a point on  $BC$  such that  $CQ = \frac{1}{3}BC$ . Prove that  $AQCP$  is a parallelogram.



27.  $PQRS$  is a parallelogram and  $\angle SPQ = 60^\circ$ . If the bisectors of  $\angle P$  and  $\angle Q$  meet at point  $A$  on  $RS$ , prove that  $A$  is the mid-point of  $RS$ .




28. In the given figure,  $K$  is the mid-point of side  $SR$  of a parallelogram  $PQRS$  such that  $\angle SPK = \angle QPK$ . Prove that  $PQ = 2QR$ .




29. Rima has a photo-frame without a photo in the shape of a triangle with sides  $a, b, c$  in length. She wants to find the perimeter of a triangle formed by joining the mid-points of the sides of the photo-frame. Find the perimeter of the triangle formed by joining the mid-points of the frame.

30. In the following figure,  $AL$  and  $CM$  are medians of  $\triangle ABC$  and  $LN \parallel CM$ . Prove that  $BN = \frac{1}{4} AB$ .




31. Two parallel lines  $l$  and  $m$  are intersected by a transversal  $p$  (see figure). Show that the quadrilateral formed by the bisectors of interior angles is a rectangle.




32.  $PQRS$  is a rhombus with  $\angle QPS = 50^\circ$ . Find  $\angle RQS$ .


33. In the given figure,  $ABCD$  is a square, side  $AB$  is produced to points  $P$  and  $Q$  in such a way that  $PA = AB = BQ$ . Prove that  $DQ = CP$ .



34. In the adjoining figure, points  $A$  and  $B$  are on the same side of a line  $m$ ,  $AD \perp m$  and  $BE \perp m$  and meet  $m$  at  $D$  and  $E$ , respectively. If  $C$  is the mid-point of  $AB$ , then prove that  $CD = CE$ .




35. In the given quadrilateral  $ABCD$ ,  $X$  and  $Y$  are points on diagonal  $AC$  such that  $AX = CY$  and  $BX \parallel DY$  is a parallelogram. Show that  $ABCD$  is a parallelogram.




## ➡ Long Answer Type Questions (LA)

36. In the given figure,  $ABCD$  is a parallelogram and  $E$  is the mid-point of  $AD$ . A line through  $D$ , drawn parallel to  $EB$ , meets  $AB$  produced at  $F$  and  $BC$  at  $L$ . Prove that (i)  $AF = 2DC$  (ii)  $DF = 2DL$



37. In  $\triangle ABC$ ,  $AB = 18$  cm,  $BC = 19$  cm and  $AC = 16$  cm.  $X, Y$  and  $Z$  are mid-points of  $AO, BO$  and  $CO$  respectively as shown in the figure. Find the perimeter of  $\triangle XYZ$ .



38. Prove that the line segment joining the mid-points of the diagonals of a trapezium is parallel to each of the parallel sides and is equal to half the difference of these sides.


39.  $ABCD$  is a parallelogram.  $AB$  and  $AD$  are produced to  $P$  and  $Q$  respectively such that  $BP = AB$  and  $DQ = AD$ . Prove that  $P, C, Q$  lie on a straight line.

40.  $P, Q, R$  and  $S$  are respectively the mid-points of sides  $AB, BC, CD$  and  $DA$  of quadrilateral  $ABCD$  in which  $AC = BD$  and  $AC \perp BD$ . Prove that  $PQRS$  is a square.

## ANSWERS

### OBJECTIVE TYPE QUESTIONS

1. (a) : Number of angles in a quadrilateral = 4.
2. (c) : Let the measure of fourth angle be  $x$ .  
Now, sum of angles of a quadrilateral =  $360^\circ$   
 $\Rightarrow 70^\circ + 120^\circ + 50^\circ + x = 360^\circ$   
 $\Rightarrow 240^\circ + x = 360^\circ \Rightarrow x = 120^\circ$
3. (b) : Here,  $X$  = Sum of angles of a triangle =  $180^\circ$ ,  
 $Y$  = Sum of angles of a quadrilateral =  $360^\circ$   
 Now,  $2X = 2 \times 180^\circ = 360^\circ = Y$   
 $\therefore 2X = Y$
4. (a) : Let the quadrilateral be  $ABCD$  in which  
 $\angle A = 90^\circ$ ,  $\angle B = 2x$ ,  $\angle C = 3x$  and  $\angle D = 4x$ .  
 Then,  $\angle A + \angle B + \angle C + \angle D = 360^\circ$   
 $\Rightarrow 90^\circ + 2x + 3x + 4x = 360^\circ$   
 $\Rightarrow 9x = 270^\circ \Rightarrow x = 30^\circ$   
 $\therefore \angle B = 60^\circ$ ,  $\angle C = 90^\circ$ ,  $\angle D = 120^\circ$   
 Hence, the largest angle is  $120^\circ$ .
5. (b) : We have,  $x + \angle A = 180^\circ$  (Linear pair)  
 $\Rightarrow x = 180^\circ - \angle A$  similarly,  $y = 180^\circ - \angle B$ ,  
 $z = 180^\circ - \angle C$ ,  $t = 180^\circ - \angle D$   
 $\Rightarrow x + y + z + t = 720^\circ - (\angle A + \angle B + \angle C + \angle D)$   
 $= 720^\circ - 360^\circ = 360^\circ$
6. (b) : In a trapezium, only one pair of opposite sides are parallel.
7. (c) : A blackboard is in the shape of a rectangle.
8. (b) : Diagonals of a rhombus are perpendicular to each other. So, the angle between them is  $90^\circ$ .
9. (c) : In a square, all the four sides are equal and all the angles are of equal measure, i.e.,  $90^\circ$ .
10. (d) : Diagonals of a kite are not equal.
11. (c) : We know, the angles of a square are bisected by the diagonals.  
 $\therefore \angle OCX = 45^\circ$   
 Also,  $\angle COD + \angle COX = 180^\circ$  (Linear pair)  
 $\Rightarrow 105^\circ + \angle COX = 180^\circ \Rightarrow \angle COX = 180^\circ - 105^\circ = 75^\circ$   
 Now, in  $\triangle COX$ , we have  
 $\angle OCX + \angle COX + \angle OXC = 180^\circ$   
 $\Rightarrow 45^\circ + 75^\circ + x = 180^\circ$   
 $\Rightarrow x = 180^\circ - 120^\circ = 60^\circ$ .
12. (c) : Let the angles of quadrilateral  $ABCD$  be  $3x$ ,  $7x$ ,  $6x$  and  $4x$  respectively.  
 $\therefore 3x + 7x + 6x + 4x = 360^\circ$   
 [Angle sum property of a quadrilateral]  
 $\Rightarrow 20x = 360^\circ$   
 $\Rightarrow x = 18^\circ$   
 $\therefore$  Angles of the quadrilateral are  
 $\angle A = 3 \times 18^\circ = 54^\circ$   
 $\angle B = 7 \times 18^\circ = 126^\circ$   
 $\angle C = 6 \times 18^\circ = 108^\circ$   
 and  $\angle D = 4 \times 18^\circ = 72^\circ$



Now, for the line segments  $AD$  and  $BC$ , with  $AB$  as transversal  $\angle A$  and  $\angle B$  are co-interior angles.

$$\text{Also, } \angle A + \angle B = 54^\circ + 126^\circ = 180^\circ$$

$$\therefore AD \parallel BC$$

Thus,  $ABCD$  is a trapezium.

13. (c) : Sum of adjacent angles of a parallelogram is  $180^\circ$ .

$$\therefore \angle A + \angle B = 180^\circ \Rightarrow 75^\circ + \angle B = 180^\circ \Rightarrow \angle B = 105^\circ$$

14. (a) :  $\angle ABC + \angle BAD = 180^\circ$

( $\because$  Sum of adjacent angles of a parallelogram is  $180^\circ$ )

$$\Rightarrow \angle ABC = 180^\circ - 70^\circ = 110^\circ$$

$$\Rightarrow \angle ABD = \angle ABC - \angle DBC = 110^\circ - 70^\circ = 40^\circ$$

Now,  $CD \parallel AB$  and  $BD$  is transversal.

$$\therefore \angle CDB = \angle ABD = 40^\circ \quad (\text{Alternate angles})$$

15. (a) : Given,  $ABCD$  is a parallelogram.

$\therefore AD \parallel BC$  and  $DF$  is a transversal.

$$\therefore \angle ADF = \angle DFC = 60^\circ \quad (\text{Alternate angles})$$

Also,  $\angle BFD + \angle DFC = 180^\circ$  (Linear pair)

$$\Rightarrow \angle BFD + 60^\circ = 180^\circ \Rightarrow \angle BFD = 180^\circ - 60^\circ = 120^\circ$$

16. (d) : Let the other two angles be  $3x$  and  $5x$ .

Now, sum of angles of a quadrilateral =  $360^\circ$ .

$$\therefore 55^\circ + 65^\circ + 3x + 5x = 360^\circ$$

$$\Rightarrow 120^\circ + 8x = 360^\circ \Rightarrow 8x = 240^\circ \Rightarrow x = 30^\circ$$

$\therefore$  Two angles are  $90^\circ$  and  $150^\circ$ .

17. (a) : Diagonals of quadrilateral bisect each other at right angle.

$\therefore$  It is a square or a rhombus.

Also, all the sides of square or rhombus are equal.

$$\therefore CD = 5 \text{ cm.}$$

18. (a) : In  $\triangle ADC$ ,

$x + y + \angle ADC = 180^\circ$  (By angle sum property of a triangle)

$$\Rightarrow \angle ADC = 180^\circ - (x + y) \quad \dots(i)$$

$$\therefore \angle ABC = \angle ADC$$

( $\because$  Opposite angles of parallelogram are equal)

$$\therefore z = 180^\circ - (x + y) \quad [\text{Using (i)}]$$

$$\Rightarrow z + x + y = 180^\circ$$

19. (a) : If a pair of opposite sides of a quadrilateral is equal and parallel, then it is a parallelogram.

20. (a) : Here,  $EF \parallel BC$  and  $F$  is mid-point of  $AC$ .

$\therefore$  By converse of mid-point theorem,  $E$  is the mid-point of  $AB$ .

$$\Rightarrow AB = 2(AE) = 2 \times 3.5 \text{ cm} = 7 \text{ cm}$$

21. (c) : Let  $ABC$  be an equilateral triangle.

$$\therefore AB = BC = AC \quad \dots(i)$$

Let  $D, E, F$  are mid-points of sides


$BC, AC, AB$  respectively.

$\therefore$  By mid-point theorem,

$$DE = \frac{1}{2} AB, EF = \frac{1}{2} BC, DF = \frac{1}{2} AC$$


$$\therefore DE = EF = DF$$

Hence,  $DEF$  is an equilateral triangle.



(From (i))

**22. (a) :** Let  $ABC$  be the triangle and  $D, E, F$  are mid-points of sides  $BC, AC, AB$  respectively.




$\therefore$  By mid-point theorem,  
 $DE \parallel AB, EF \parallel BC, DF \parallel AC$   
 $\therefore DEAF, BDEF, FDCE$  are all parallelograms.  
 Now,  $DE$  is the diagonal of parallelogram  $FDCE$   
 $\therefore \triangle DEC \cong \triangle EDF$

Similarly,  $\triangle FAE \cong \triangle EDF$

and  $\triangle BFD \cong \triangle EDF$

Hence, all four triangles are congruent.

**23. (d) :** Given,  $M$  and  $N$  are respectively mid-points of non-parallel sides  $PS$  and  $QR$  of trapezium  $PQRS$ .



Join  $RM$  and produce it to meet  $QP$  produced at  $X$ .

In  $\triangle SMR$  and  $\triangle PMX$ ,

$\angle SMR = \angle PMX$  (Vertically opposite angles)

$\angle SRM = \angle PXM$

( $\because$  Alternate angles as,  $SR \parallel QX$  and  $XR$  is transversal)  
 $SM = PM$  ( $\because M$  is mid-point of  $PS$ )

$\therefore \triangle SMR \cong \triangle PMX$  (By AAS congruence rule)

$\Rightarrow MR = MX$  and  $SR = PX$  (By C.P.C.T.)

Now, in  $\triangle RXQ$ ,  $M$  is the mid-point of  $XR$ , as  $XM = MR$  and  $N$  is the mid-point of  $RQ$ .

$\therefore$  By mid-point theorem,  $MN \parallel XQ$  and  $MN = \frac{1}{2} XQ$

$\Rightarrow MN \parallel PQ$  and  $MN = \frac{1}{2} (XP + PQ) = \frac{1}{2} (SR + PQ)$   
 ( $\because SR = XP$ )

Hence,  $MN \parallel PQ$  and  $MN = \frac{1}{2} (SR + PQ)$

**24. (b) :** In  $\triangle CDB$ , we have  $CD = CB$

[ $\because$  adjacent sides of rhombus are equal]

$\Rightarrow \angle CBD = \angle CDB = x$

In  $\triangle BCD$ ,  $\angle BCD = 70^\circ$

and  $\angle CDB + \angle CBD + \angle DCB = 180^\circ$

$\Rightarrow x + x + 70^\circ = 180^\circ \Rightarrow x = 55^\circ$

$\Rightarrow \angle CDB = 55^\circ$

**25. (b) :** Since, adjacent angles of a parallelogram are supplementary.

So,  $2x + 25^\circ + 3x - 5^\circ = 180^\circ$

$\Rightarrow 5x = 160^\circ \Rightarrow x = 32$

**26. (a) :** Let the three angles  $\angle T, \angle A$  and  $\angle R$  be  $5x, 3x$  and  $7x$  respectively.

$\therefore \angle S + \angle T + \angle A + \angle R = 360^\circ$

$\Rightarrow 120^\circ + 5x + 3x + 7x = 360^\circ$

$\Rightarrow 15x = 240^\circ \Rightarrow x = 16^\circ$

$\therefore \angle R = 7 \times 16 = 112^\circ$

**27. (b) :** Since  $ABCD$  is a trapezium.

$\therefore x + 20^\circ + 2x + 10^\circ = 180^\circ$

(Sum of measure of interior angles is  $180^\circ$ )

$\Rightarrow 3x + 30 = 180^\circ \Rightarrow x = 50^\circ$

and  $y + 92^\circ = 180^\circ \Rightarrow y = 88^\circ$

**28. (a) :** Given  $\angle A + \angle C = 2(\angle B + \angle D)$

$\Rightarrow 140^\circ + \angle C = 2\angle B + 2 \times 60^\circ$

$\Rightarrow 2\angle B - \angle C = 20^\circ$

...(i)

Also,  $\angle A + \angle B + \angle C + \angle D = 360^\circ$

$\Rightarrow 140^\circ + \angle B + \angle C + 60^\circ = 360^\circ$

$\Rightarrow \angle B + \angle C = 160^\circ$

...(ii)

Using (i) and (ii), we get  $\angle B = 60^\circ$

**29. (c) :** Let the smallest angle be  $\angle A = x^\circ$ ,

and other angle be  $\angle B = (2x - 24)^\circ$

$\therefore \angle A + \angle B = 180^\circ$

$\Rightarrow x + 2x - 24 = 180$


$\Rightarrow 3x = 204 \Rightarrow x = 68$

$\therefore \angle A = 68^\circ$

and  $\angle B = (2x - 24)^\circ = (2 \times 68 - 24)^\circ = 112^\circ$

Since, opposite angles of a parallelogram are equal.

So,  $\angle A = \angle C = 68^\circ, \angle B = \angle D = 112^\circ$



**30. (c) :** Let the measures of the angles be  $2x, 4x, 5x$  and  $7x$ .

$\Rightarrow 2x + 4x + 5x + 7x = 360^\circ$

(Angle sum property)


$\Rightarrow 18x = 360^\circ \Rightarrow x = 20^\circ$

$\therefore \angle A = 40^\circ, \angle B = 80^\circ, \angle C = 100^\circ, \angle D = 140^\circ$

As  $\angle A + \angle D = 180^\circ$  and  $\angle B + \angle C = 180^\circ$

$\Rightarrow CD \parallel AB$

$\therefore ABCD$  is a trapezium.




**31. (c) :** In  $\triangle PQR$ ,  $A$  and  $B$  are mid-points of  $PQ$  and  $PR$  respectively.

$\therefore AB \parallel QR$  [By mid-point theorem]

$\therefore \angle AQR = \angle PAB$

[Corresponding angles]

$\therefore \angle PQR = \angle PAB = 60^\circ$



**32. (c) :** We have,  $\angle ADC + b = 180^\circ$

[Linear pair]

$\Rightarrow \angle ADC = 180^\circ - b$

...(i)

Also,  $\angle ABC + a = 180^\circ$

[Linear pair]

$\Rightarrow \angle ABC = 180^\circ - a$

...(ii)

In quadrilateral  $ABCD$ , we have

$\angle ABC + \angle BCD + \angle ADC + \angle BAD = 360^\circ$

[By angle sum property of a quadrilateral]

$\Rightarrow 180^\circ - a + y + 180^\circ - b + x = 360^\circ$  [Using (i) and (ii)]

$\Rightarrow 360^\circ - a - b + x + y = 360^\circ$

$\Rightarrow x + y = a + b$

**33. (b) :** Given,  $PQRS$  is a parallelogram.

$\therefore SR \parallel PQ$  and  $SR = PQ$

...(i)

But,  $QT = PQ$  (Given)

...(ii)

From (i) and (ii), we have  $SR = PQ = QT$

In  $\triangle SRO$  and  $\triangle TQO$

$\angle RSO = \angle QTO$

(Alternate angles)

$SR = QT$

(Proved above)

$\angle SRO = \angle TQO$

(Alternate angles)

$\therefore \triangle SRO \cong \triangle TQO$

(By ASA congruency criteria)

$\Rightarrow RO = OQ$

(By C.P.C.T.)

**34. (b) :** If consecutive sides of a parallelogram are equal, then it is necessarily a rhombus.



35. (d) : Let  $ABC$  be right angled triangle and  $\angle ABC = 90^\circ$ .

Let  $D, E, F$  are mid-points of sides  $BC, AC$  and  $AB$  respectively.

$\therefore EF \parallel BD$  and  $BF \parallel DE$

(By mid-point theorem)

$\Rightarrow BDEF$  is a parallelogram.

$\therefore \angle FED = \angle FBD = 90^\circ$

( $\because$  Opposite angles of a parallelogram are equal)

$\therefore DEF$  is right angled triangle.

36. (b) : A parallelogram can be formed by joining the mid points of sides of quadrilateral.

37. (b) : As  $P$  and  $Q$  are mid points of  $AB$  and  $AD$  respectively.

$\therefore PQ = \frac{1}{2}BD$  ... (1)

and  $PQ \parallel BD$  [By midpoint theorem]

38. (a) : As,  $R$  and  $S$  are mid points of  $CD$  and  $BC$  respectively.

$\therefore RS \parallel BD$  and  $RS = \frac{1}{2}BD$  i.e.,  $BD = 2RS$  ... (2)

39. (b) : From (1) and (2),  $RS = PQ = \frac{1}{2}BD$

40. (a) : Perimeter of quadrilateral  $PQRS$   
 $= PQ + QR + RS + SP$

41. (d) : All the conditions given in options (a), (b) and (c) are necessary for  $ABCD$  to be a quadrilateral.

42. (c) : In a parallelogram, diagonal can't bisect each other.

43. (a) :  $\angle A = \angle C \Rightarrow \frac{1}{2}\angle A = \frac{1}{2}\angle C$

$\Rightarrow \angle YAX = \angle YCX$

Also,  $\angle AYC + \angle YCX = 180^\circ$

[ $\because CX \parallel AY$ ]

$\therefore \angle AYC + \angle YAX = 180^\circ$

So,  $AX \parallel CY$  ( $\because$  Interior angles on the same side of the transversal are supplementary)

44. (b) : As  $ABCD$  is a parallelogram.

$\therefore \angle A = \angle C = 63^\circ$

(Opposite angles of a parallelogram are equal)

Also,  $AEFG$  is a parallelogram.

$\therefore \angle A + \angle G = 180^\circ$  (Adjacent angles are supplementary)

$\therefore \angle G = 180^\circ - 63^\circ = 117^\circ$

45. (c) : Let the angles be  $3x, 5x, 5x$  and  $7x$ .

Now,  $3x + 5x + 5x + 7x = 360^\circ$

$\Rightarrow 20x = 360^\circ \Rightarrow x = 18^\circ$

$\therefore$  All angles are  $54^\circ, 90^\circ, 90^\circ, 126^\circ$

46. (b) : In  $\triangle ADE$  and  $\triangle CFE$ , we have

$AE = CE$

(Given)


$DE = FE$

(Given)

$\angle AED = \angle CEF$

(Vertically opposite angles)

$\therefore \triangle ADE \cong \triangle CFE$  (By SAS congruency criterion)



47. (b) :  $\angle EFC = \angle EDA$

(By CPCT)

48. (a) :  $\angle ECF = \angle EAD$

(By CPCT)

49. (d) :  $CF = AD$

(By CPCT)

50. (c) :  $CF \parallel BD$


( $\because \angle ECF = \angle EAD$ )

51. (c) : In quadrilateral  $ABXC$ , we have  
 $AD = DX$  [Given]

$BD = DC$  [Since  $AD$  is median]

So, diagonals  $AX$  and  $BC$  bisect each other but not at right angles.

Therefore,  $ABXC$  is a parallelogram.




52. (a) : Clearly, statement-II is true.

Now, in  $\triangle ADC$ ,  $Q$  is the mid-point of  $AC$  such that  $PQ \parallel AD$ .

$\therefore P$  is the mid-point of  $DC$ .

[By converse of mid-point theorem]

$\Rightarrow DP = PC$




53. (a) : Since, opposite angles of a parallelogram are equal. Therefore,  $3x - 2 = 50 - x \Rightarrow x = 13$ .

So, angles are  $(3 \times 13 - 2)^\circ = 37^\circ$  and  $(50 - 13)^\circ = 37^\circ$ .

54. (a) : Since, diagonals of a square bisect each other at right angles.

$\therefore \angle AOB = 90^\circ$



55. (b) : In  $\triangle ABC$ ,  $E$  and  $F$  are midpoint of the sides  $AC$  and  $AB$  respectively.


$\therefore FE \parallel BC$  [By mid-point theorem]

Now, in  $\triangle ABP$ ,  $F$  is mid-point of  $AB$  and

$FQ \parallel BP$

$\Rightarrow Q$  is mid-point of  $AP$

$\Rightarrow AQ = QP$ .



## SUBJECTIVE TYPE QUESTIONS

1. We know that consecutive interior angles of a parallelogram are supplementary.

$\therefore (x + 60^\circ) + (2x + 30^\circ) = 180^\circ$

$\therefore 3x + 90^\circ = 180^\circ \Rightarrow 3x = 90^\circ \Rightarrow x = 30^\circ$

Thus, two consecutive angles are  $(30^\circ + 60^\circ)$ ,  $(2 \times 30^\circ + 30^\circ)$  i.e.,  $90^\circ$  and  $90^\circ$ .

Hence, the special name of the given parallelogram is rectangle.

2.  $\angle PQR = \angle PSR = 125^\circ$

( $\because$  Opposite angles of a parallelogram are equal)

Now,  $\angle PQR + \angle RQT = 180^\circ$

(Linear pair)

$\Rightarrow 125^\circ + \angle RQT = 180^\circ \Rightarrow \angle RQT = 55^\circ$

3. No.

$\therefore$  Sum of the angles  $= 110^\circ + 80^\circ + 70^\circ + 95^\circ$   
 $= 355^\circ \neq 360^\circ$

Thus, the given angles cannot be the angles of a quadrilateral.

4. Yes,  $QL = LR$

As, opposite sides of a parallelogram are equal.

$\therefore$  In parallelogram  $QLMN$ ,  $QL = NM$  ... (i)

In parallelogram  $NLRM$ ,  $NM = LR$  ... (ii)

From (i) and (ii),  $QL = LR$

5. Since,  $\angle A + \angle B = 180^\circ$

[Co-interior angles]

$$\Rightarrow \angle B = 180^\circ - 78^\circ = 102^\circ$$

Now,  $\angle B = \angle D = 102^\circ$

and,  $\angle A = \angle C = 78^\circ$

[ $\therefore$  opposite angles of a parallelogram are equal]

6. We have,  $\angle A + \angle B = 180^\circ$  [Co-interior angles]

$$\Rightarrow 60^\circ + \angle ABD + 55^\circ = 180^\circ \Rightarrow \angle ABD = 65^\circ$$

Also,  $\angle ABD = \angle CDB$

[Alternate interior angles are equal]

$$\therefore \angle CDB = \angle ABD = 65^\circ$$

We have,  $\angle ADB = \angle DBC$

[Alternate interior angles are equal]

$$\Rightarrow \angle ADB = 55^\circ$$

7. We have,  $AB = AC \Rightarrow \angle BCA = \angle B$

Now,  $\angle CAD = \angle B + \angle BCA$  [Exterior angle property]

$$\Rightarrow 2\angle CAP = 2\angle BCA \quad [\because AP \text{ is the bisector of } \angle CAD]$$

$$\Rightarrow \angle CAP = \angle BCA \Rightarrow AP \parallel BC$$

Also,  $AB \parallel CP$

[Given]

Hence,  $ABCP$  is a parallelogram.

8. If one angle of a rhombus is a right angle, then it is necessarily a square.

9. Since a rhombus is a parallelogram.

$\therefore$  Its opposite angles are equal.

$$\Rightarrow \angle A = \angle C$$

$$\therefore \angle C = 60^\circ \quad [\because \angle A = 60^\circ \text{ (Given)}]$$

$$\text{Now, required sum} = \angle A + \angle C = 60^\circ + 60^\circ = 120^\circ$$

10. We have given, a trapezium  $ABCD$ , whose parallel sides are  $AB$  and  $DC$ .

Since,  $AB \parallel CD$  and  $AD$  is a transversal.

$$\therefore \angle A + \angle D = 180^\circ \quad [\text{Angles on same side of transversal}]$$

$$\Rightarrow \angle D = 180^\circ - \angle A = 180^\circ - 45^\circ = 135^\circ$$

Similarly,  $\angle C = 135^\circ$

11. In  $\triangle COD$ , we have

$$\angle COD + \angle 1 + \angle 2 = 180^\circ$$

$$\Rightarrow \angle COD = 180^\circ - (\angle 1 + \angle 2)$$





$$\Rightarrow \angle COD = 180^\circ - \left( \frac{1}{2}\angle C + \frac{1}{2}\angle D \right)$$

$$\Rightarrow \angle COD = 180^\circ - \frac{1}{2}(\angle C + \angle D)$$

$$\Rightarrow \angle COD = 180^\circ - \frac{1}{2}\{360^\circ - (\angle A + \angle B)\}$$

$$[\because \angle A + \angle B + \angle C + \angle D = 360^\circ]$$

$$\Rightarrow \angle COD = \frac{1}{2}(\angle A + \angle B)$$



12. In  $\triangle BCD$ , we have

$$\angle BDC + \angle DCB + \angle CBD = 180^\circ$$

[Angle sum property of a triangle]

$$\Rightarrow 5a + 9a + 4a = 180^\circ$$

$$\Rightarrow 18a = 180^\circ \Rightarrow a = 10^\circ$$

$$\therefore \angle C = 9 \times 10^\circ = 90^\circ$$

Since, opposite angles of a parallelogram are equal

$$\text{Therefore, } \angle A = \angle C \Rightarrow \angle A = 90^\circ$$

13. True. Given,  $ABCD$  is a quadrilateral whose diagonals bisect each other. Then, it should be a parallelogram.

Also,  $\angle A$  and  $\angle B$  are adjacent angles of parallelogram  $ABCD$ . So, their sum should be  $180^\circ$ .

$$\text{Now, } \angle A + \angle B = 45^\circ + 135^\circ = 180^\circ$$

14. Since,  $D$  and  $E$  are the mid-point of sides  $AB$  and  $AC$  respectively.

$$\therefore AD = \frac{1}{2}AB \text{ and } AE = \frac{1}{2}AC$$

$$\text{By mid-point theorem, } DE = \frac{1}{2}BC$$

$$\therefore AD + AE + DE = \frac{1}{2}(AB + AC + BC)$$

$$\text{Perimeter of } \triangle ADE = \frac{1}{2} \times \text{perimeter of } \triangle ABC$$

$$= \frac{1}{2} \times 35 \text{ cm} = 17.5 \text{ cm}$$

Hence, the perimeter of  $\triangle ADE$  is 17.5 cm.

15. In  $\triangle ABC$ ,  $DE \parallel AB$  and  $AD$  is the median.

So,  $D$  is the mid-point of  $BC$ .

By converse of mid-point theorem,

$E$  is the mid-point of  $AC$ .

Hence,  $BE$  is median.

16. We have,

$$MN = \frac{1}{2}BC, MP = \frac{1}{2}AC \text{ and } NP = \frac{1}{2}AB$$

[By midpoint theorem]

$$\Rightarrow BC = 6 \text{ cm, } AC = 5 \text{ cm}$$

and  $AB = 7 \text{ cm}$ .

The value of  $(BC + AC) - AB$

$$= (6 + 5) - 7 = 4 \text{ cm.}$$

17.  $ABC$  is an isosceles triangle with  $AB = AC$  and  $D, E$  and  $F$  as the mid-points of sides  $BC, CA$  and  $AB$  respectively.  $AD$  intersects  $FE$  at  $O$ . Join  $DE$  and  $DF$ .




Since,  $D, E$  and  $F$  are mid-points of sides  $BC, AC$  and  $AB$  respectively.

$$\therefore DE \parallel AB \text{ and } DE = \frac{1}{2}AB \quad [\text{By mid-point theorem}]$$

$$\text{Also, } DF \parallel AC \text{ and } DF = \frac{1}{2}AC$$

But,  $AB = AC$

[Given]



$$\Rightarrow \frac{1}{2}AB = \frac{1}{2}AC$$

$$\Rightarrow DE = DF \quad \dots(i)$$

$$\text{Now, } DE = \frac{1}{2}AB \Rightarrow DE = AF \quad \dots(ii)$$

$$\text{and, } DF = \frac{1}{2}AC \Rightarrow DF = AE \quad \dots(iii)$$

From (i), (ii) and (iii), we have

$DE = AE = AF = DF \Rightarrow DEAF$  is a rhombus.

Since, diagonals of a rhombus bisect each other at right angles.

$\therefore AD \perp FE$  and  $AD$  is bisected by  $FE$ .

18. Here,  $\angle ABE + \angle EBF = 90^\circ$

$$\Rightarrow 30^\circ + \angle EBF = 90^\circ$$

$$\Rightarrow \angle EBF = 60^\circ \quad \dots(i)$$

and  $\angle BFE + \angle CFE = 180^\circ$  [Linear pair]

$$\Rightarrow \angle BFE + 144^\circ = 180^\circ$$

$$\Rightarrow \angle BFE = 180^\circ - 144^\circ = 36^\circ \quad \dots(ii)$$

Now, in  $\triangle BEF$ ,

$$\angle EBF + \angle BFE + \angle BEF = 180^\circ \quad (\text{Angle sum property})$$

$$\Rightarrow 60^\circ + 36^\circ + \angle BEF = 180^\circ \quad [\text{Using (i) and (ii)}]$$

$$\Rightarrow \angle BEF = 180^\circ - 96^\circ = 84^\circ$$

19. Let  $ABCD$  be a parallelogram with  $AB$  and  $DC$  as longer sides and  $AD$  and  $BC$  as shorter sides.

Now,  $AB = DC = 9.5$  cm [Opposite sides of a parallelogram are equal and longer side = 9.5 cm (Given)]

Let  $AD = BC = x$

$$\text{Now, } AB + BC + CD + DA = 30 \quad [\text{Perimeter} = 30 \text{ cm (Given)}]$$

$$\Rightarrow 9.5 + x + 9.5 + x = 30$$

$$\Rightarrow 2x = 30 - 19 = 11 \Rightarrow x = 5.5 \text{ cm}$$

$\therefore$  Length of shorter side = 5.5 cm

20. Since,  $ABCD$  is a parallelogram.

$$\therefore \angle A = \angle C$$

$$\Rightarrow (3x - 20)^\circ = (x + 40)^\circ$$

$$\Rightarrow 3x - x = 40 + 20$$

$$\Rightarrow 2x = 60 \Rightarrow x = 30$$

Also,  $\angle A + \angle B = 180^\circ$

$$\Rightarrow (3x - 20)^\circ + (y + 15)^\circ = 180^\circ$$

$$\Rightarrow 3x + y = 185 \Rightarrow y = 185 - 90 = 95$$

$$\therefore x + y = 30 + 95 = 125$$

21.  $\angle SPQ = \angle QRS = 2x$

( $\because$  Opposite angles of a parallelogram are equal)

In  $\triangle PSQ$ ,  $\angle PSQ + \angle PQS + \angle SPQ = 180^\circ$

$$\Rightarrow 4x + 4x + 2x = 180^\circ$$

$$\Rightarrow 10x = 180^\circ$$

$$\Rightarrow x = 18^\circ$$

Now,  $\angle PSR = \angle PQR$

( $\because$  Opposite angles of a parallelogram are equal)

$$\Rightarrow 4x + \angle QSR = 4x + \angle SQR$$

$$\Rightarrow \angle QSR = \angle SQR \quad \dots(i)$$

$$\text{In } \triangle SRQ, \angle SRQ + \angle RSQ + \angle SQR = 180^\circ$$

$$\Rightarrow 2 \times 18^\circ + 2 \angle RSQ = 180^\circ \quad [\text{From (i)}]$$

$$\Rightarrow 2 \angle RSQ = 180^\circ - 36^\circ = 144^\circ \Rightarrow \angle RSQ = 72^\circ$$

Hence,  $\angle P = \angle R = 2 \times 18^\circ = 36^\circ$ ,

$$\angle Q = \angle S = 4x + 72^\circ = 4 \times 18^\circ + 72^\circ = 144^\circ$$

22. We have,  $AB = BC$  and have to prove that  $DE = EF$ . Now, trapezium  $ACFD$  is divided into two triangles namely  $\triangle ACF$  and  $\triangle AFD$ .

In  $\triangle ACF$ ,  $AB = BC \Rightarrow B$  is mid-point of  $AC$

and  $BG \parallel CF$

[ $\because m \parallel n$ ]

So,  $G$  is the mid-point of  $AF$ .

[By converse of mid-point theorem]

Now, in  $\triangle AFD$ ,  $G$  is the mid-point of  $AF$ .

and  $GE \parallel AD$

[ $\because m \parallel l$ ]

$\therefore E$  is the mid-point of  $FD$ .

[By converse of mid-point theorem]

$$\Rightarrow DE = EF$$

$\therefore l, m$  and  $n$  cut off equal intercepts on  $q$  also.

23. Let  $ABCD$  be the rhombus and greater diagonal  $AC$  be  $x$  cm.

$$\therefore \text{Smaller diagonal, } BD = \frac{1}{3}AC = \frac{x}{3} \text{ cm}$$

Since diagonals of rhombus are perpendicular bisector of each other.


$$\therefore OA = \frac{x}{2} \text{ cm and } OB = \frac{x}{6} \text{ cm}$$

In  $\triangle AOB$ , we have

$$AB^2 = OA^2 + OB^2$$

$$\Rightarrow 10^2 = \left(\frac{x}{2}\right)^2 + \left(\frac{x}{6}\right)^2$$

$$\Rightarrow 100 = \frac{x^2}{4} + \frac{x^2}{36} \Rightarrow 100 = \frac{10}{36}x^2 \Rightarrow x = 6\sqrt{10} \text{ cm}$$



24. Let  $D, E$  and  $F$  be the mid-points of sides  $BC, CA$  and  $AB$  respectively.

In  $\triangle ABC$ ,  $F$  and  $E$  are mid-points of  $AB$  and  $AC$ .

$$\therefore FE \parallel BC \text{ and } FE = \frac{1}{2}BC$$

$$\therefore FE \parallel BD \text{ and } FE = BD$$

$\therefore FEDB$  is a parallelogram.

Similarly,  $CDFE$  and  $AFDE$  are also parallelograms.

$$\therefore \angle B = \angle DEF, \angle C = \angle DFE \text{ and } \angle FDE = \angle A$$

$$\Rightarrow \angle DEF = 60^\circ, \angle DFE = 70^\circ \text{ and } \angle FDE = 50^\circ$$

25. Suppose  $AC$  and  $BD$  intersect at  $O$ .

$$\text{Then, } OC = \frac{1}{2}AC$$

$$\text{Now, } CQ = \frac{1}{4}AC \quad [\text{Given}]$$

$$\Rightarrow CQ = \frac{1}{2}OC$$





In  $\triangle COD$ ,  $P$  and  $Q$  are the midpoints of  $DC$  and  $OC$  respectively.

$$\therefore PQ \parallel DO$$

[By mid-point theorem]

Also, in  $\triangle COB$ ,  $Q$  is the midpoint of  $OC$  and  $QR \parallel OB \therefore R$  is the midpoint of  $BC$ .

[By converse of mid-point theorem]



26.  $\therefore ABCD$  is parallelogram.

$\Rightarrow AD = BC$  and  $AD \parallel BC$

$\Rightarrow \frac{1}{3}AD = \frac{1}{3}BC$  and  $AD \parallel BC$

$\Rightarrow AP = CQ$  and  $AP \parallel CQ$

Thus,  $APCQ$  is a quadrilateral such that one pair of opposite sides  $AP$  and  $CQ$  are parallel and equal.

Hence,  $APCQ$  is a parallelogram.

27.  $\angle P + \angle Q = 180^\circ$

(Adjacent angles of parallelogram)

$\Rightarrow 60^\circ + \angle Q = 180^\circ \Rightarrow \angle Q = 120^\circ$

Since,  $PA$  and  $QA$  are bisectors of angles  $P$  and  $Q$

$\therefore \angle SPA = \angle APQ = \frac{1}{2}\angle P = \frac{1}{2} \times 60^\circ = 30^\circ$

And  $\angle RQA = \angle AQP = \frac{1}{2}\angle Q = \frac{1}{2} \times 120^\circ = 60^\circ$

Now,  $SR \parallel PQ$  and  $AP$  is transversal.

$\therefore \angle SAP = \angle APQ = 30^\circ$  [Alternate interior angles]

In  $\triangle ASP$ , we have

$\angle SAP = \angle APS = 30^\circ$

$\Rightarrow SP = AS$

(Sides opposite to equal angles are equal) ... (i)

Similarly,  $QR = AR$  ... (ii)

But,  $QR = SP$  [Opposite sides of parallelogram] ... (iii)

From (i), (ii) and (iii), we have  $AS = AR$

$\Rightarrow A$  is the mid-point of  $SR$ .

28. We have,  $\angle SPK = \angle QPK$  ... (i)

Now,  $PQ \parallel RS$  and  $PK$  is a transversal

$\therefore \angle SKP = \angle QPK$  [Alternate angles] ... (ii)

From (i) and (ii),  $\angle SPK = \angle SKP$

$\Rightarrow PS = SK$  ... (iii)

( $\because$  Sides opposite to equal angles are equal)

But  $K$  is the mid-point of  $SR$ .

$\therefore SK = KR$  ... (iv)

$PS = QR$  (Opposite sides of parallelogram are equal) ... (v)

From (iii) and (v),  $SK = PS = QR$

Also,  $PQ = SR = SK + KR = 2SK$  [From (i)]  
 $= 2QR$

29. Let the photo-frame

be  $ABC$  such that  $BC = a$ ,

$CA = b$  and  $AB = c$  and the

mid-points of  $AB$ ,  $BC$  and  $CA$

are  $D$ ,  $E$  and  $F$  respectively.

We have to determine the

perimeter of  $\triangle DEF$ .


In  $\triangle ABC$ ,  $DF$  is the line-segment joining the mid-points of sides  $AB$  and  $AC$ .

By mid-point theorem,  $DF \parallel BC$  and  $DF = \frac{BC}{2} = \frac{a}{2}$

Similarly,  $DE = \frac{AC}{2} = \frac{b}{2}$  and  $EF = \frac{AB}{2} = \frac{c}{2}$

$\therefore$  Perimeter of  $\triangle DEF = DF + DE + EF$

$$= \frac{a}{2} + \frac{b}{2} + \frac{c}{2} = \frac{a+b+c}{2}$$



30. We have,  $AL$  and  $CM$  are medians of  $\triangle ABC$ , i.e.,  $L$  and  $M$  are the mid-points of  $BC$  and  $AB$  respectively.

$\therefore LC = BL = \frac{1}{2}BC$  and  $BM = AM = \frac{1}{2}AB$  ... (i)

In  $\triangle BMC$ ,  $L$  is the mid-point of  $BC$  and  $LN \parallel CM$ . So, by converse of mid-point theorem,  $N$  is mid-point of  $BM$ .

i.e.,  $BN = NM = \frac{1}{2}BM$  ... (ii)

From (i) and (ii), we get

$$BN = \frac{1}{2} \left( \frac{1}{2}AB \right) \Rightarrow BN = \frac{1}{4}AB$$

31. It is given that  $l \parallel m$  and transversal  $p$  intersects them at points  $A$  and  $C$  respectively.

The bisectors of  $\angle PAC$  and  $\angle ACQ$  intersect at  $B$  and bisectors of  $\angle ACR$  and  $\angle SAC$  intersect at  $D$ .

Now,  $\angle PAC = \angle ACR$

[Alternate angles as  $l \parallel m$  and  $p$  is a transversal]

So,  $\frac{1}{2}\angle PAC = \frac{1}{2}\angle ACR$

$\Rightarrow \angle BAC = \angle ACD$

These form a pair of alternate angles for lines  $AB$  and  $DC$  with  $AC$  as transversal and they are equal also.

So,  $AB \parallel DC$

Similarly,  $BC \parallel AD$

Therefore, quadrilateral  $ABCD$  is a parallelogram.

Also,  $\angle PAC + \angle CAS = 180^\circ$  [Linear pair]

So,  $\frac{1}{2}\angle PAC + \frac{1}{2}\angle CAS = \frac{1}{2} \times 180^\circ = 90^\circ$

$\Rightarrow \angle BAC + \angle CAD = 90^\circ \Rightarrow \angle BAD = 90^\circ$

So,  $ABCD$  is a parallelogram in which one angle is  $90^\circ$ .

Therefore,  $ABCD$  is a rectangle.

32. Since a rhombus satisfies all the properties of a parallelogram.

$\therefore \angle QPS = \angle QRS$

[Opposite angles of a parallelogram]

$\Rightarrow \angle QRS = 50^\circ$

[ $\because \angle QPS = 50^\circ$  (Given)]

$\therefore$  Diagonals of a rhombus bisect the opposite angles.

$\therefore \angle ORQ = \frac{1}{2}\angle QRS \Rightarrow \angle ORQ = 25^\circ$

Now, in  $\triangle ORQ$ , we have

$\angle OQR + \angle ORQ + \angle ROQ = 180^\circ$

$\Rightarrow \angle OQR + 25^\circ + 90^\circ = 180^\circ$

[ $\because$  Diagonals of a rhombus are perpendicular

to each other  $\Rightarrow \angle ROQ = 90^\circ$ ]

$\Rightarrow \angle OQR = 180^\circ - 115^\circ = 65^\circ$

$\therefore \angle RQS = 65^\circ$

33. Since,  $ABCD$  is a square.

$\therefore AB = BC = CD = DA$

Also,  $PA = AB = BQ$


$\therefore AB = BC = CD = DA = PA = BQ$

In  $\triangle PDA$  and  $\triangle QCB$ ,  $PA = BQ$

$AD = BC$

(Given)

(Sides of a square)



$$\begin{aligned}
 \angle A &= \angle B && \text{(Each } 90^\circ) \\
 \Delta PDA &\cong \Delta QCB && \text{(By SAS congruency rule)} \\
 \Rightarrow PD &= QC && \text{(By C.P.C.T.)} \dots(i) \\
 \angle PDA &= \angle QCB && \text{(By C.P.C.T.)} \dots(ii) \\
 \text{Now, } \angle PDC &= \angle PDA + \angle ADC && \\
 &= \angle PDA + 90^\circ && \dots(iii) \\
 \angle QCD &= \angle QCB + \angle BCD && \\
 &= \angle QCB + 90^\circ && \dots(iv)
 \end{aligned}$$

From (ii), (iii) and (iv), we have  $\angle PDC = \angle QCD$   
 Now, in  $\Delta PDC$  and  $\Delta QCD$ ,  $PD = QC$  (Given)  
 $DC = DC$  (Common)  
 $\angle PDC = \angle QCD$  (Proved above)  
 $\Delta PDC \cong \Delta QCD$  (By SAS congruency rule)  
 $\therefore DQ = CP$  [By C.P.C.T.]

34. We have, C is the mid-point of AB

$$\therefore AC = BC.$$

Draw,  $CM \perp m$  and join AE.

We have,  $AD \perp m$ ,

$$CM \perp m \text{ and } BE \perp m.$$

$$\therefore AD \parallel CM \parallel BE$$

In  $\Delta ABE$ ,  $CG \parallel BE$  [ $\because CM \parallel BE$ ]

and C is the mid-point of AB.

Thus, by converse of mid-point theorem, G is the mid-point of AE.

In  $\Delta ADE$ , G is the mid-point of AE and  $GM \parallel AD$ .

$$[\because CM \parallel AD]$$

Thus, by converse of mid-point theorem, M is mid-point of DE.

In  $\Delta CMD$  and  $\Delta CME$ ,

$$DM = EM \quad (\because M \text{ is the mid-point of } DE)$$

$$\angle CMD = \angle CME = 90^\circ \quad (\because CM \perp m)$$

$$CM = CM \quad \text{(Common)}$$

$$\therefore \Delta CMD \cong \Delta CME \quad \text{(By SAS congruency rule)}$$

$$\text{So, } CD = CE \quad \text{(By C.P.C.T.)}$$

35. Since BXYD is a parallelogram.

$$\therefore XO = YO \quad \dots(i)$$

$$\text{and } DO = BO \quad \dots(ii)$$

[ $\because$  Diagonals of a parallelogram bisect each other]

$$\text{Also, } AX = CY \quad \text{(Given)} \quad \dots(iii)$$

$$\text{Adding (i) and (iii), we have } XO + AX = YO + CY$$

$$\Rightarrow AO = CO \quad \dots(iv)$$

From (ii) and (iv), we have

$$AO = CO \text{ and } DO = BO$$

Thus, ABCD is a parallelogram, because diagonals AC and BD bisect each other at O.

36. (i) As  $EB \parallel DF \Rightarrow EB \parallel DL$  and  $ED \parallel BL$ .

Therefore, EBLD is a parallelogram.

$$\therefore BL = ED = \frac{1}{2} AD = \frac{1}{2} BC = CL \quad \dots(i)$$

$$[\because ABCD \text{ is a parallelogram } \therefore AD = BC]$$

Now, in  $\Delta DCL$  and  $\Delta FBL$ , we have

$$CL = BL \quad \text{[from (i)]}$$

$$\angle DLC = \angle FLB \quad \text{(Vertically opposite angles)}$$

$$\angle DCL = \angle FBL \quad \text{(Alternate angles)}$$

$$\therefore \Delta DCL \cong \Delta FBL \quad \text{(By ASA congruency criteria)}$$

$$\Rightarrow CD = BF \text{ and } DL = FL \quad \text{(By C.P.C.T.)}$$

$$\text{Now, } BF = DC = AB \quad \dots(ii)$$

$$\Rightarrow 2AB = 2DC \Rightarrow AB + AB = 2DC$$

$$\Rightarrow AB + BF = 2DC$$

[Using (ii)]

$$\Rightarrow AF = 2DC$$

$$(ii) \therefore DL = FL \Rightarrow DF = 2DL$$

37. Here, in  $\Delta ABC$ ,  $AB = 18$  cm,  $BC = 19$  cm,

$AC = 16$  cm.

In  $\Delta AOB$ , X and Y are the mid-points of AO and BO.

$\therefore$  By mid-point theorem, we have

$$XY = \frac{1}{2} AB = \frac{1}{2} \times 18 \text{ cm} = 9 \text{ cm}$$

In  $\Delta BOC$ , Y and Z are the mid-points of BO and CO.

$\therefore$  By mid-point theorem, we have

$$YZ = \frac{1}{2} BC = \frac{1}{2} \times 19 \text{ cm} = 9.5 \text{ cm}$$

And, in  $\Delta COA$ , Z and X are the mid-points of CO and AO.

$\therefore$  By mid-point theorem, we have

$$\therefore ZX = \frac{1}{2} AC = \frac{1}{2} \times 16 \text{ cm} = 8 \text{ cm}$$

Hence, the perimeter of  $\Delta XYZ = 9 + 9.5 + 8 = 26.5$  cm

38. Let, trapezium ABCD in which,  $AB \parallel DC$  and P and Q are the mid-points of its diagonals AC and BD respectively.

We have to prove (i)  $PQ \parallel AB$  and  $PQ \parallel DC$

$$(ii) \quad PQ = \frac{1}{2} (AB - DC)$$

Join D and P and produce DP to meet AB at R.

(i) Since  $AB \parallel DC$  and transversal AC cuts them at A and C respectively.

$$\therefore \angle 1 = \angle 2 \quad \text{(Alternate angles)} \dots(1)$$

In  $\Delta APR$  and  $\Delta CPD$ ,

$$\angle 1 = \angle 2 \quad \text{(From (1))}$$

$$AP = CP \quad (\because P \text{ is the mid-point of } AC)$$

$$\angle 3 = \angle 4 \quad \text{(Vertically opposite angles)}$$

$$\therefore \Delta APR \cong \Delta CPD \quad \text{(By ASA congruency rule)}$$

$$\Rightarrow AR = DC \text{ and } PR = DP \quad \text{(By C.P.C.T.)}$$

In  $\Delta DRB$ , P and Q are the mid-points of side DR and DB respectively.

$$\therefore PQ \parallel RB \quad \text{(By mid-point theorem)}$$

$$\Rightarrow PQ \parallel AB \quad (\because RB \text{ is a part of } AB)$$



$$\Rightarrow PQ \parallel AB \text{ and } PQ \parallel DC \quad (\because AB \parallel DC)$$

(ii) In  $\Delta DRB$ , P and Q are the mid-points of side DR and DB respectively.

$$\therefore PQ = \frac{1}{2} RB \quad \text{(By mid-point theorem)}$$

$$\Rightarrow PQ = \frac{1}{2} (AB - AR) \Rightarrow PQ = \frac{1}{2} (AB - DC)$$

[From part (i),  $AR = DC$ ]





39. CP and CQ are joined.

∴ ABCD is a parallelogram.

So, BC = AD, AB = DC

[Opposite sides of parallelogram]  
and  $\angle ABC = \angle ADC$

[Opposite angles of parallelogram]

∴ Their supplementary angles are equal

So,  $\angle PBC = \angle CDQ$

In  $\triangle PBC$  and  $\triangle CDQ$ , we have

BC = DQ [BC = AD and AD = DQ (Given)]

BP = DC [AB = DC and AB = BP (given)]

$\angle PBC = \angle CDQ$  [Proved above]

∴  $\triangle PBC \cong \triangle CDQ$  [By SAS congruency]

⇒  $\angle BPC = \angle DCQ$  and  $\angle BCP = \angle DQC$  [By C.P.C.T.]

Again,  $\angle BCD = \angle PBC$  [since, AP || DC]

Now,  $\angle BCP + \angle BCD + \angle DCQ$

$= \angle BCP + \angle PBC + \angle BPC = 2 \text{ right angles}$

i.e.,  $\angle PCQ$  is a straight angle.

i.e., P, C, Q lie on a straight line.


40. In quadrilateral ABCD,  $AC \perp BD$  and  $AC = BD$ .

In  $\triangle ADC$ , S and R are the mid-points of the sides AD and DC respectively.

∴  $SR \parallel AC$  and  $SR = \frac{1}{2} AC$  ... (i)

[By mid-point theorem]

In  $\triangle ABC$ , P and Q are the mid-points of AB and BC respectively.



∴  $PQ \parallel AC$  and  $PQ = \frac{1}{2} AC$  ... (ii)

[By mid-point theorem]

From (i) and (ii),  $PQ \parallel SR$

and  $PQ = SR = \frac{1}{2} AC$  ... (iii)

Similarly, in  $\triangle ABD$ ,

$SP \parallel BD$  and  $SP = \frac{1}{2} BD$  [By mid-point theorem]

∴  $SP = \frac{1}{2} AC$  [ $\because AC = BD$ ] ... (iv)

Now in  $\triangle BCD$ ,  $RQ \parallel BD$  and  $RQ = \frac{1}{2} BD$

[By mid-point theorem]

∴  $RQ = \frac{1}{2} AC$  [ $\because BD = AC$ ] ... (v)

From (iv) and (v),  $SP = RQ = \frac{1}{2} AC$  ... (vi)

From (iii) and (vi),  $PQ = SR = SP = RQ$  ... (vii)

∴ All four sides are equal.

Now, in quadrilateral OERF,

$OE \parallel FR$  and  $OF \parallel ER$

∴  $\angle EOF = \angle ERF = 90^\circ$

[ $\because AC \perp DB$ ]

∴  $\angle QRS = 90^\circ$

... (viii)

From (vii) and (viii), we get  
 $PQRS$  is a square.

